

DOI 10.26724/2079-8334-2025-1-91-212-219

UDC 614.4; 616-08

M.F. Amirova, E.E. Huseynova, N.V. Melikova, F.I. Mammadova, U.H. Azizova, S.R. Guliyeva
Azerbaijan Medical University, Baku, Azerbaijan

**BIOLOGICALLY SUBSTANTIATED, SAFE,
AND CORRECTIVE REMEDIES FOR SARS-COV-2 REGIMEN**

e-mail: gerayelmira@gmail.com

The paper seeks to find a novel effective treatment way for Severe Acute Respiratory Syndrome COVID-2 (SARS-CoV-2) without harm to the organism in general. Taking into account the information on the mechanism of SARS-CoV-2 development, during treatment, it is advisable to pay attention to inhibition of viral replication; reduction hypoxia; elimination drug poisoning and toxic, allergic consequences; add antioxidant therapy to prevent formation of reactive oxygen species (ROS) at recovery phase and hook up anti-inflammatory therapy. This goal could be achieved by enhancing antiviral defense, administering antioxidants, and activating antioxidant enzymes according to their action mechanisms and therapeutic implications. For example, using superoxide dismutase mimics metalloporphyrins, metalocorroles, Mn biliverdins, Mn cyclic polyamines, Mn salens, metal oxides/salts and inducers of antioxidant protection. It is also possible to increase the antioxidant enzymes genes expression or the activity of the enzyme itself. Patients taking antibiotics for SARS-CoV-2 were more susceptible to exacerbations due to opportunistic infections, so alternatives to antibiotics with high efficacy and well-absorbed antioxidant properties are more appropriate. In this regard, vitamin C, N-acetylcysteine, and glycine, as well as melatonin, quercetin, astaxanthin, glutathione, fat-soluble vitamins, polyphenols, unsaturated fatty acids, and a number of other substances with high anti-inflammatory effects exhibit positive effect.

Key words: antioxidants, antiviral activity, cytokine storm, inflammation, oxidative stress, SARS-CoV-2, vitamins.

М.Ф. Амірова, Е.Е. Гусейнова, Н.В. Мелікова, Ф.І. Мамедова, У.Х. Азізова, С.Р. Гулієва

**БІОЛОГІЧНО ОБГРУНТОВАНІ, БЕЗПЕЧНІ І КОРИГУЮЧІ ЗАСОБИ
ДЛЯ ЛІКУВАННЯ SARS-COV-2**

У статті розглянуто новий науково-обґрунтований, клінічно перевірений ефективний спосіб лікування SARS-CoV-2 без шкоди для організму. З огляду на механізм розвитку SARS-CoV-2 при лікуванні доцільно приділяти увагу наступним моментам: гальмуванню реплікації вірусу; зменшенню гіпоксії; усуненню лікарського отруєння та токсичних, алергічних наслідків; додати антиоксидантну терапію для запобігання утворенню активних форм кисню (ROS) на етапі відновлення та підключити протизапальну терапію. Ця мета може бути досягнута шляхом посилення противірусного захисту, застосуванню антиоксидантів та активації антиоксидантних ферментів відповідно до механізмів їх дії та терапевтичних наслідків. Наприклад, використання супероксиддисмутази імітує металопорфірини, металококоролі, Mn білівердини, Mn циклічні поліаміни, Mn саліни, оксиди/солі металів та індуктори антиоксидантного захисту. Також можна збільшити експресію генів антиоксидантних ферментів або активність самого ферменту. Пацієнти, які приймають антибіотики для лікування SARS-CoV-2, були більш схильні до загострень через опортуністичні інфекції, тому більш підходящими є альтернативи антибіотикам з високою ефективністю та добре засвоюваними антиоксидантними властивостями. У цьому відношенні позитивний вплив мають вітамін С, N-ацетилцистеїн і гліцин, а також мелатонін, кверцетин, астаксантин, глутатіон, жиророзчинні вітаміни, поліфеноли, ненасичені жирні кислоти та ряд інших речовин з високою протизапальною дією.

Ключові слова: антиоксиданти, цитокіновий шторм, запалення, окислювальний стрес, SARS-CoV-2, вітаміни, противірусна дія.

Previous outbreaks of acute respiratory syndrome are known to have occurred in 2002 and 2012, but they were characterized by relatively low prevalence vs 2019's [18]. This recent, SARS-CoV-2, was characterized by the highest virulence and susceptibility. The high mortality rate, as well as the serious economic and social disadvantages, has made the study of the mechanisms of the disease and therapeutic methods the most pressing problem in medicine. However, the exact pathogenesis of this infection has not yet been identified, but the collected data suggest that the virus is a factor that causes polyorganic pathology and damages not only the lungs, but also the vascular wall, hemostasis. The virus is also a major contributor to comorbid pathology, characterized by an increase in reactive oxygen species (ROS) and disruption of the immune system. Virus-induced ROS and redox imbalance exacerbate the inflammatory response to SARS-CoV infection, ergo the mitochondria, the source of ROS, are closely monitored. The pathogenesis of coronavirus has been closely studied for nearly 20 years, although their effective prophylaxis and therapeutic methods have not yet been developed. There are a number of reasons for this, some of which include the ability of the virus to mutate intensively, high virulence of strains, features of the pathogenetic course, rapid spread. To date, the interaction of viral activity with oxidative stress, the effect of these relationships on cells and tissues are the leading factors in the development of acute stress syndrome, and the role of antioxidant deficiency in the genesis of infection has also been elucidated.

The active forms of oxygen formed during metabolism are of great importance, as they act as messengers in the cell. It is known that ROS-dependent kinases can stimulate inflammatory signals in the cell by increasing the expression and transcription of inflammatory genes. Activation of antimicrobial cells, including neutrophils and macrophages, and the formation of inflammatory cytokines are mainly dependent on ROS. Optimal levels of ROS in the body are provided by the enzymatic and non-enzymatic antioxidant system (AOS). When the antioxidant defenses are weakened, a condition called oxidative stress occurs. Numerous studies have shown that increased ROS accompanies all infectious respiratory viral infections, as well as HIV and hepatitis [8, 9]. The activation of a number of pathological processes that cause complications for the body occurs along with the increase in ROS, and the development of coronavirus infection does not go beyond this rule. Coronaviruses (Latin Coronaviridae) are a class of about 40 zoonotic RNA-containing viruses that are widespread in the human population and among animals. There are 5 main types of coronaviruses, two of which (α - and β -viruses) infect humans. The first human coronavirus (HCoV-B814) was first detected in 1965 in a patient with an acute respiratory virus infection. At the beginning of the 21st century, coronaviruses were known as veterinary pathogens and did not pose a threat to humans. Coronavirus infections have been found to be moderate, with mild symptoms of the upper respiratory tract, but β -coronavirus epidemics in recent decades have overturned previous views. The SARS-CoV coronavirus, which caused China's severe acute respiratory syndrome in 2002-2003, and the Middle East coronavirus (MERS-CoV) in 2012, killed more than 10,000 people; Mortality in SARS-CoV was 10 % and in MERS-CoV it was 37 %. In late 2019, a new type of coronavirus, identified by the WHO as SARS-CoV-2 and causing a new severe acute respiratory syndrome (COVID-19), was identified.

The purpose of the study was to investigate the biologically substantiated, safe, and corrective remedies for SARS-COV-2 regimen.

Material and methods. SARS-related data were collected from PubMed, WHO Global Research Database, ClinicalTrials.gov, Scopus journals, Web of Science (WoS) high-impact journals, and Google Scholar diverse scholarly articles. We concentrated our analysis on data from 2019 to 2023, a period marked by the emergence of SARS-CoV-2 and an unprecedented intensification of research into therapeutic approaches and treatments for coronavirus-related infections. This timeframe captures the global scientific response to the pandemic, including advancements in antiviral therapies, immunomodulatory strategies, and the evaluation of both traditional and novel treatment modalities. To capture a comprehensive range of studies addressing both traditional and contemporary approaches to managing SARS and retrieve relevant data, we employed targeted keywords reflecting our research focus, including “SARS and Immune response”, “Antiviral therapies for SARS”, “Biologically safe treatments for SARS”, “Phytotherapy for SARS” “Natural remedies for SARS”.

Current understanding and emerging treatment strategies for SARS-CoV-2 cellular invasion mechanisms, and therapeutic approaches.

Recent genetic research has shown that the virus originally appeared in bats, but whether the bats transmitted the virus to themselves is still being investigated. Epidemic SARS-CoV-2 bats in humans are closely related to two coronavirus types, bat-SL-CoVZC45 and bat-SL-CoVZXC21, which manifest themselves in the β -CoV / bat / Yunnan / RaTG 13/2013 sequence.

The SARS-CoV-2 genome repeats the MERS-CoV genome by 50 %, the SARS-CoV-1 genome – by 79 %, and the BtRs-CoV genome – by 88 %. Analysis of 86 complete and incomplete genomes of SARS-CoV-2 revealed numerous mutations in its encoded and unencoded areas, indicating that the new strain of coronavirus has extensive genetic material and its ability to evolve rapidly.

The virus is a circular pleomorphic pathogen with a lipid coating of 80–229 nm; On its surface are three pathogenic structural proteins. On the outside of the virus are crown-shaped glycoprotein protrusions (spike S-protein) designed to bind to the surface of the target cell. It has been found that spike proteins have the ability to hide from innate immunity by undergoing conformational changes. The spike protein of the SARS-CoV-2 virus binds to target cells through angiotensin converting enzyme-2 (ACE2), which forms active angiotensin. ACE2 is mainly expressed in cells of the gastrointestinal tract, kidneys, blood vessels, heart and lungs. The SARS virus can also bind to the CD147 cell receptor, also known as BASIGIN. As a result of the virus attaching to the cell, the spike protein is broken down into S1 and S2 subunits by proteolysis of the cell via type 2 transmembrane serine protease (transmembrane serine 2 protease, TMPRSS2). S1 then binds to the subunit ACE2, then the S1-RBD PD-ACE2 complex dissociates, the hydrophobic S2-FP peptide (fusion peptide) is released from S2, and activating endocytosis, the virus enters the target cell. Then the virus permanent mechanism of action is activated: viral RNA is translated through the target cell's organelles, resulting in the formation of structural proteins necessary for virus development,

which also form new SARS-CoV virions, and continue to damage new target cells. ACE2 and CD147 receptors sensitive to SARS-CoV-2 and TMPRSS2 are located on the surface of the upper respiratory tract and gastrointestinal epithelial cells, which are the entry routes of infection. The products of the virus interaction with the target cell are recognized by specific NOD-receptors [2] involved in the formation of the polyprotein complex, the inflammasome [21, 26].

RNA viruses induce the accumulation and activation of NLRP3-inflammasome in the early stages of the disease, this inflammasome plays a leading role in the formation of immunity against the virus. The virus metabolites produce ROS, which damage mitochondria and stimulate the release of DNA from mitochondria. Heat shock protein A1L (Heat Shock Protein A1L, HSPA1L) increases viral replication in the host cell; in SARS-CoV infection the rate of DNA methylation decreases because COVID reduces the activity of DNA methyl-transferases in epithelial cells of lung tissue [22]. This leads to hyperexpression of the A1L heat shock protein during SARS-CoV-2, and consequently the spread of the virus [22]. In contrast to strains with low virulence, SARS-CoV-2 has the ability to damage type I and II alveocytes, as well as endothelial cells by passing into the lower respiratory tract. This process results in the transcription expression and of anti-inflammatory cytokines. During the cytokine secretion phase, the alveolar epithelium undergoes pyroptosis, the resulting products are absorbed by granulocytes and tissue macrophages [5]. In this case, neutrophils and cytotoxic T cells, along with the formed cyto- and chemokines, join the process for protection lung tissue from the virus. During the fight against the virus, edema develops at the site of injury resulting in severe pulmonary pneumonia that causes acute respiratory distress syndrome and fibrosis. The age factor aggravates the pathological process, i.e. these changes listed are more likely to manifest themselves more pronounced in the elderly [22]. When the body immune response to the SARS-CoV-2 virus is insufficient, the virus enters the bloodstream and can damage other organs and cells that have receptors on their surfaces: intestines, kidneys, esophagus, heart, blood vessels, brain, bladder, etc.

It is also thought, that SARS-CoV-2 disease is a type of general viral vasculitis, and damage to the lungs in this disease is one different type of angiogenic edema. The distinctive feature of SARS-CoV-2 is the widespread involvement of the immune system during development of this pathogenesis, which leads to the elimination of SARS-CoV-2 in the early stages, but in the late stages of severe inflammation stimulates severe or even fatal dysfunction of some organs. Thus, due to the stimulation of endogenous immunomodulators synthesis by COVID-19, severe forms of “cytokine storm” occur, which results in the loss of control of the inflammatory process and a significant deficiency of organs and systems [15]. In the early stages of the disease, the virus nsp1 and rp6 proteins inhibit the formation of interferon. Macrophages entering the site of inflammation continue to produce chemoattractants for single-nucleated cells (mononuclear cells), and thus their concentration increases rapidly, which stimulates the inflammatory process to the next stage, the “cytokine storm”. At this stage, levels of inflammatory cytokines and hemattractants continue to increase dramatically, as interleukins, monocyte hematopoietic protein MCP-1, macrophage inflammatory protein MIP-1a, TGF, CCL2, CXCL10, TNF (CXCL9) increase sharply [24].

In severe cases of SARS-CoV-2 infection, there are significant changes in the acute phase of the blood, and at this stage, inflammation markers such as C-reactive protein, ferritin, ceruloplasmin, blood clotting factors, indicative serum enzymes indicate a lack of multiple organs [23, 30]. In about 82 % of cases of peripheral blood, leukopenia, thrombocytopenia and loss of eosinophils with lymphopenia are observed. For this reason, SARS infection is characterized by the appearance of significant immunopathological elements. Coronaviruses can damage both the upper and lower parts of the human respiratory system. Most of the clinical signs do not differ in multiple types of viruses, in severe cases the disease results in bronchitis, pneumonia and acute respiratory distress syndrome.

Viral respiratory infections stimulate the inflammatory process and promote the development of pathophysiological processes against the background of an excess of active forms of cytokines and oxygen and/or nitrogen. The main generator of ROS is mitochondria, the SARS-CoV-3b protein of the COVID virus and the non-structural protein 10 (nsp10) can change the course of processes in mitochondria [16, 31, 32]. SARS-CoV 3b may enter mitochondria, while nsp10 may interact specifically with the NADH 4L subunit and cytochromoxidase II. It has also been shown that genes encoding mitochondrial DNA in peripheral blood mononuclear cells, as well as genes sensitive to oxidative stress such as peroxiredoxin 1, activate the ferritin heavy-chain polypeptide gene. Oxidative stress increases the expression of 2D type of anti-inflammatory phospholipase A2, which reduces antiviral immunity. Interestingly, in humans, phospholipase A2 2D is naturally activated as ages [20, 29]

In humans, there is also protein kinase, which is activated by mitogens from the destructive effects of the external environment, such as oxidative stress, accumulation of DNA damage, carcinogens and viruses. Phosphorylated activated mitogenic protein kinases were found in all SARS-CoV-infected cells. A number of studies confirm that there is a direct link between oxidative stress and congenital immune damage, and that this mechanism is one of the main causes of lung tissue damage in SARS-CoV-2. Thus, those who have a deletion of the *ncf1* gene that stimulates the formation of ROS and do not have Toll-like receptor-4 (TLR4) which activates innate immunity, have a natural resistance to respiratory viruses, including SARS COVID virus. This is explained by the fact that organisms with such changes in the body do not produce oxidized 1-palmitoyl-2-arachidoinyl-phosphatidyl choline, which stimulates the formation of cytokines by macrophages and activates damaging mechanisms. It is also known that 3CLpro protease of SARS-CoV induces apoptosis in human promonocytes by increasing the formation of ROS. Occurrence of oxidative stress and activation of transcription factor NF-KB can lead to severe lung damage [17]. In vivo oxidative stress, 3CLpro activates NF- κ B-factor but inhibits protein-1-dependent transcription, so ROS-induced transmission of 3CLpro-stimulated NF-KB signal may have played a leading role in the pathophysiology of SARS-CoV-2 infection [1, 15]. This is because the activation of the phosphatidylinositol-3-kinase/protein kinase pathway by various viruses during latent or chronic infections allows infected cells to avoid apoptosis, which in turn contributes to the spread of the virus in the body.

The severity of SARS-CoV infection increases with age [25]. This is explained by age-related changes in the immune system, the accumulation of oxidized products, and, conversely, the weakening of antioxidant defenses, and the consequent accumulation of mass OAF due to the violation of the oxidation-reduction balance. In the late stages, the redox-sensitive factor transcription factor NF-KB is activated, which is accompanied by the induction of inflammatory cytokines (IL-6, TNF- α), an increase in adhesion molecules. Oxidative stress triggered by SARS-CoV and NF-KB transmission by Toll receptors can further increase the reactivity of target cells, resulting in severe lung damage. Age-related changes are more pronounced in SARS-CoV infections, especially in pandemics, where the number of deaths from SARS-CoV-2 is less than 0.2 % in a population under 60 and close to 10 % over 80 years of age. Hypertension, diabetes, obesity, and other such type diseases increase the risk of death in this age group to 5 %. At present, coronavirus infection unequivocally belongs to the group of pathologies caused by free radical reactions. Thus, after SARS-CoV enters the body, infiltration of macrophages, monocytes and neutrophils into the lung tissue begins, and due to the presence of the virus in the body, they start to actively form ROS. In fact, this mechanism, that is the conversion of oxygen into active forms serves for defense cells, it is self-defense, because ROS has a lethal effect on the virus & virus-infected cells. FAD-dependent NADPH-oxidase of mitochondria rapidly increases the number of ROS by transferring electrons to oxygen. This condition is called "oxidative stress" because it leads to a tenfold increase in oxygen reactive forms. Naturally, ROS primarily serves to remove damaged, weak, functionally unnecessary cells in the injury zone. However, in the process of cleaning the damaged area, the activity of some relatively healthy cells which are closely adjacent to it, is impaired. In order to control the process, the body begins to produce anti-inflammatory cytokines, and while they increase in the blood, circulating neutrophils can adhere to the site of injury and damage the pulmonary endothelium. Proteases of neutrophils & ROS that break down lung proteins and increase so rapidly, that they are thought to be key elements in SARS-CoV infection, which damage lung tissue. The aggressive reaction of neutrophils, as we have noted, is primarily directed against the infected cell, so the cytotoxic effect of neutrophils weakens as the number of cells infected with the virus decreases. However, when the process gets out of control and the virus is widespread in the body, acute respiratory distress syndrome occurs. As the damaged cells are destroyed by neutrophils, the patient's blood and kidney secretions begin to increase the products of DNA degradation, as well as malon dialdehyde, keto- and hydroxycholesterol, formed by the peroxide oxidation of lipids. Depending on the degree of damage to the lungs, the breakdown products of protein (carbonyls) in the blood and urine increase. Nitric oxide is added to the process of cleansing the body of the virus, which is accompanied by an increase in nitroguanosine. Deficiency of the antioxidant system leads to the fatal outcome of SARS-CoV infection.

Thus, the increase in the expression of the heavy ferritin chain (FTH1) in SARS-CoV infection [21–24] indicates the need to strengthen the antioxidant system, because ferritin stores iron and prevents its formation in ROS by Fenton reactions, thus vitamin crease in other antioxidant iron-binding proteins that limit the rate of ROS formation. Of course, in the progressive stage of SARS-CoV-2 infection, when the virus spreads rapidly in the body, the need for prooxidants, rather than antioxidants, increases. This is because ROS eliminates damaged cells along with the virus and prevents the spread of infection. For this reason, at this stage the concentration of proteins that play an antioxidant role in the body and regulate iron

metabolism – haptoglobin, ferritin, transferrin and ceruloplasmin – decreases in the blood. Naturally, the lack of the above-mentioned proteins, which prevent the cascade reactions of ROS formation during the recovery phase and are actively involved in the maintenance of innate immunity, can result in unpleasant complications for the body. Taking into account the information on the mechanism of development of SARS-CoV-2 in the literature on homeostasis disorders, it is advisable to pay attention to the inhibition of viral replication; reduction of hypoxia; elimination of drug poisoning and toxic, allergic consequences; antioxidant therapy to prevent the formation of ROS during the recovery phase; anti-inflammatory therapy. This goal could be achieved through the administration of antioxidants with anti-viral properties, as well as the use of antiviral drugs with antioxidant effects [7, 26]. However, the literature suggests that antioxidant therapy alone is less effective in SARS-CoV-2 infection, so treatment should focus on the use of drugs used in previous respiratory infections. For example, using superoxide dismutase mimics (metalloporphyrins, metallocorroles, Mn biliverdins, Mn cyclic polyamines, Mn salens, and metal oxides and salts) [19] and inducers of antioxidant protection, scientists have proposed substances that stop the development of the disease and increase the effect of each other in lower respiratory tract infections. To achieve this, it is possible to increase the expression of genes of antioxidant enzymes or the activity of the enzyme itself.

Antioxidants lead to a reduction in virus-induced nuclear factor NF- κ B [11] and the interferon regulatory factor-3 (IRF-3) transcription factors [12, 13], as well as cytokines and chemokines. It has also been suggested, that a “cytokine storm” must be prevented in order to control systemic inflammation [14]. Despite the antiviral activity of cytokines, their overproduction during this infection has a stronger destructive effect on lung tissue than the virus itself. Interferons belong to the antiviral group, but interferons themselves belong to the cytokine class, which leads to additional activation of immune cells, which overshadows the use of these drugs in the case of SARS-CoV-2. For this reason, there is currently an active search for alternatives to interferon, and these drugs include antioxidants that provide enzymatic and non-enzymatic protection of the body [11]. One of the factors that increase the body's non-enzymatic antioxidant defenses is vitamin C, which can have an anti-inflammatory effect by rapidly reducing the formation of ROS. Due to its antiviral activity, vitamin C also calms the “cytokine storm” in the body against the virus, and from this point of view, it is more suitable than interferon [10]. Interestingly, vitamin C can be used both alone and in combination with other drugs that have a synergistic effect with it. Thus, vitamin C with sulforaphane has a positive effect in the treatment of acute inflammatory diseases requiring artificial ventilation of the lungs [3, 13, 6]. Vitamin C is currently actively used in the treatment of SARS-CoV-2 in China, and this antioxidant has even been included in the ClinicalTrials.gov protocols. Thiol antioxidants, such as glutathione, are also involved in the cell's non-enzymatic defense system because they break down peroxide radicals during viral respiratory disease, weakening apoptosis and inhibiting viral replication [33]. In vitro analysis of the antiviral effects of 5 antioxidant components such as tocopherol, thiamine, pantothenic acid, pyridoxine, biotin and glutathione showed that most glutathione and pyridoxine were active, while thiamine, biotin and tocopherol also had a fairly high inhibitory effect. Unsaturated fatty acids omega-3 and omega-6, as well as vitamins E, A and D, and metal ions with variable valence are significantly higher in antioxidant and anti-inflammatory effects, so they are recommended for use in the treatment of SARS-CoV patients.

Recent studies have found a strong link between vitamin D deficiency and COVID-19 infection in the elderly population in different countries [25]. In addition to the mucolytic effect of N-acetyl-L-cysteine, it can prevent the induction of anti-apoptosis and anti-inflammatory IL-6, IL-8 cytokines, and has a strong antiviral effect. Vitamin C and N-acetyl-cysteine are currently the most widely used antioxidants in lung damage [26, 27]. The antiviral effects of some flavonoids are directly related to their inhibition of 3C-like proteases. Thus, the preparation of quercetin from the group of natural flavonoids (Fig.1) was found to have a negative effect on the synthesis of viral RNA and the level of cytokines in the blood [34]. Studies have also shown significant antiviral activity of resveratrol and catrexin drugs from the group of natural polyphenols. These compounds have been shown to be effective against viral replication as they have delayed expression of viral proteins, ergo these small polar molecules with a fused ring structure, most of which belong to the polyphenols group, will be used in the future as SARS-CoV-2 antiviral drugs. Anti-inflammatory and antioxidant, anti-coronaviral effects of taurine, carnosine and 4-hydroxyproline from animal origine has been found [28, 29]. Studies have shown that the enzyme branch of the antioxidant defense system can restore the damaged regulatory mechanisms of the respiratory system, as well as significantly reduce viral titers under the influence of antioxidant enzymes. For example, the main antioxidant enzyme that inactivates superoxide radicals is superoxide dismutase and catalase, which inactivates hydrogen peroxide.




Fig.1. Some classes of substances preventing SARS-CoV-2 from entry host cell, replication, and spread [34].

Melatonin, which is rapidly absorbed, also has significant antioxidant properties. There are data in the literature on the inhibitory effect of melatonin on necrotic pyroptosis – programmed cell death caused by NLRP3-inflammation inhibition during SARS-CoV [30, 31]. Studies have shown that melatonin and α -lipoic acid can significantly reduce the damaging effects of antiviral drugs. The antioxidant properties of melatonin are indirectly due to the activation of antioxidant enzymes in the body, which allows to increase the body's overall defenses. It is known that melatonin deficiency occurs with age [24, 28], and this feature should be taken into account in the treatment of elderly SARS-CoV patients. The anti-stress effects of melatonin have also been confirmed, which has a positive effect in the COVID pandemic.

Another antioxidant, astaxanthin, has high anti-inflammatory activity because it reduces C-reactive protein, the effects of interleukin-1 and a number of other inflammatory mediators of the respiratory tract [10].

Recently, information has been published on the use of a well-known antiseptic, leukomethylene (a reduced form of methylene oxide), approved by the World Health Organization for use in the treatment of a wide range of diseases, for the treatment of SARS-CoV-2 patients [4]. This substance along with vitamin C, N-acetylcysteine and α -lipoic acid, has a significant antioxidant effect, prevents the spread of viral RNA, leading to a weakening of inflammation and hypoxia.

In the treatment of SARS-CoV-2 infection, there is a need to eliminate toxic side effects arising by the use of antiviral drugs, with antioxidant drugs. It is known that the protection of biological structures from ROS is carried out by different antioxidant mechanisms, which are realized through different regulatory pathways. Therefore, from a pathogenetic point of view, in order to ensure synergism and drug accumulation, antioxidant drugs should be prescribed not in isolated form, but in a complex way, that stimulates both enzymatic and non-enzymatic defense, and this aspect is already reflected in the treatment of SARS-CoV-2 patients. Thus, according to the adopted protocol, prevention and treatment of SARS-CoV-2 patients is carried out through a “vitamin cocktail” containing vitamins D and C, methionine, zinc, quercetin. Such a combination of these components, which provide antioxidant protection, is expedient from both a prophylactic and therapeutic point of view.

The development of vaccines can be expected, but it is a time-consuming and costly process, additionally, vaccine injection should be avoided in some group of patients, such as in diseases accompanied by thrombosis, hypertension and so on, because their vaccination leads to death or, at least – to undesirable complications.

The rise in uncontrolled SARS-CoV-2 infections and deaths is prompting scientists around the world to think of new strategies that could reduce the secretion of anti-inflammatory cytokines (“cytokine storms”). Loss of control of the inflammatory response can lead to even more serious consequences than the effects of the virus, and often results in a number of vital conditions, including respiratory failure, which can be fatal. Therefore, one of the main goals of the current pandemic is to quell the “cytokine storm” that occurs as a consequence of oxidative stress. Drugs currently used for anti-cytokine therapy may have

hepatotoxic effects [6]. Substances used as an alternative to them should be highly effective and have well-absorbed antioxidant properties. It is known that in the other respiratory viral infections vitamin C, N-acetylcysteine and glycine, as well as melatonin, quercetin, astaxanthin, glutathione, fat-soluble vitamins, polyphenols, unsaturated fatty acids, and a number of other substances with high anti-inflammatory effects have positive effect; these substances used as an alternative to toxic drugs show effective antioxidant properties and are well-absorbed, ergo they can be the drugs of choice in SARS-CoV-2 treatment.

Conclusions

1. Considering SARS-CoV virus mutation rate and worsening health by vaccination in patients with hypertension and weakened immune systems, a preferable approach to treating and preventing SARS-CoV-2 could involve enhancing the body's defense mechanisms with supplements like vitamins D and C, methionine, zinc, and quercetin.

2. Vitamin C, glutathione, and pyridoxine pronounced therapeutic effects can be applied in SARS-CoV management due to their antioxidant properties.

3. Thiamine, biotin, and tocopherol demonstrate a noticeable inhibitory effect on the virus.

4. Omega-3 and omega-6 fatty acids, along with vitamins E, A, D, and transition metal ions, exhibit even stronger antioxidant and anti-inflammatory effects than thiamine, biotin, and tocopherol.

5. N-acetyl-L-cysteine has significant antiviral properties due to regulation of anti-apoptotic and pro-inflammatory cytokines IL-6 and IL-8.

6. Flavonoids inhibit 3S-like proteases, thereby exerting antiviral effects.

7. Taurine, carnosine, and 4-hydroxyproline of animal origin possess anti-inflammatory, antioxidant, and antiviral properties favorable in SARS-CoV.

8. The antioxidant astaxanthin reduces C-reactive protein levels, the effects of interleukin-1, and other inflammatory mediators, providing a strong anti-inflammatory effect.

9. Melatonin inhibits necrotic pyroptosis caused by the suppression of NLRP3 inflammasome activation and, in combination with α -lipoic acid, significantly reduces the damaging effect of antiviral drugs.

10. Leukomethylene, in conjunction with vitamin C, N-acetylcysteine, and α -lipoic acid, exhibits strong antioxidant properties, reduces viral RNA spread, and mitigates inflammation and hypoxia.

References

1. Amirova, MF, Azizova, GI. SARS-CoV-2 Virus Transmission Mechanism in the Human Body. *Pakistan journal of biochemistry and molecular biology*. 2021;4(1-2): 14–20.
2. Avdonin PP, Rybakova EY, Trufanov SK, Avdonin PV. SARS-CoV-2 Receptors and Their Involvement in Cell Infection. *Biochem (Mosc) Suppl Ser A Membr Cell Biol*. 2023;17(1):1-11. doi: 10.1134/S1990747822060034.
3. Bagheri Hosseiniabadi M, Khanjani N, Norouzi P, Faghihi-Zarandi A, Darban-Sarokhalil D, Khoramrooz SS, et al. The Effects of Antioxidant Vitamins on Proinflammatory Cytokines and Some Biochemical Parameters of Power Plant Workers: A Double-Blind Randomized Controlled Clinical Trial. *Bioelectromagnetics*. 2021 Jan;42(1):18-26. doi: 10.1002/bem.22294.
4. Buckley LF, Wohlford GF, Ting C. Role for Anti-Cytokine Therapies in Severe Coronavirus Disease 2020. *Crit Care Explor*. 2020;2(8):e0178. Published 2020 Aug 10. doi:10.1097/CCE.0000000000000178.
5. Carcaterra M, Caruso C. Alveolar epithelial cell type II as main target of SARS-CoV-2 virus and COVID-19 development via NF-Kb pathway deregulation: A physio-pathological theory. *Med Hypotheses*. 2021 Jan;146:110412.
6. Chaolin H, Yeming W, Xingwang L, Lili R, Jianping Z, Yi H et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020; 395(10223):497 – 5061.
7. Chaudhary P, Janmeda P, Docea AO, Yeskalyeva B, Abdull Razis AF, Modu B, et al. Oxidative stress, free radicals and antioxidants: potential crosstalk in the pathophysiology of human diseases. *Front Chem*. 2023 May 10;11:1158198. doi: 10.3389/fchem.2023.1158198.
8. Costela-Ruiz VJ, Illescas-Montes R, Puerta-Puerta JM, Ruiz C, Melguizo-Rodríguez L. SARS-CoV-2 infection: The role of cytokines in COVID-19 disease. *Cytokine Growth Factor Rev*. 2020;54:62-75.
9. Dabholkar N, Gorantla S, Dubey SK, Alexander A, Taliyan R, Singhvi G. Repurposing methylene blue in the management of COVID-19: Mechanistic aspects and clinical investigations. *Biomed Pharmacother*. 2021 Oct;142:112023. doi: 10.1016/j.biopha.2021.112023.
10. DE Flora S, Balansky R, LA Maestra S. Antioxidants and COVID-19. *J Prev Med Hyg*. 2021 Jun 5;62(1 Suppl 3):E34-E45. doi: 10.15167/2421-4248/jpmh2021.62.1S3.1895.
11. De Melo AF, Homem-de-Mello M. High-dose intravenous vitamin C may help in cytokine storm in severe SARS-CoV-2 infection. *Crit Care*. 2020;24(1):500. Published 2020 Aug 13. doi:10.1186/s13054-020-03228-3.
12. Forcados GE, Muhammad A, Oladipo OO. Metabolic Implications of Oxidative Stress and Inflammatory Process in SARS-CoV-2 Pathogenesis: Therapeutic Potential of Natural Antioxidants REVIEW article. *Front. Cell. Infect. Microbiol.*, 26 May 2021. <https://doi.org/10.3389/fcimb.2021.654813>.
13. Gasparello J, D'Aversa E, Papi C, Gambari L, Grigolo B, Borgatti M, et al. Sulforaphane inhibits the expression of interleukin-6 and interleukin-8 induced in bronchial epithelial IB3-1 cells by exposure to the SARS-CoV-2 Spike protein. *Phytomedicine*. 2021 Jul;87:153583. doi: 10.1016/j.phymed.2021.153583.
14. Hiroyuki N, Masaaki M. Neurobiology of COVID-19-Associated Psychosis/Schizophrenia: Implication of Epidermal Growth Factor Receptor Signaling. *Neuropsychopharmacology Reports*. 2025; 45(1): e12520.

15. Jafarzadeh A, Chauhan P, Saha B, Jafarzadeh S, Nemati M. Contribution of monocytes and macrophages to the local tissue inflammation and cytokine storm in COVID-19: Lessons from SARS and MERS, and potential therapeutic interventions. *Life Sci*. 2020;257:118102.
16. Kabekkodu Sh, Chakrabarty S, Jayaram P, Mallya S, Thangaraj K, Singh KK, et al. Severe acute respiratory syndrome coronaviruses contributing to mitochondrial dysfunction: Implications for post-COVID complications. *Mitochondrion*. 2023; 69:43-56, ISSN 1567-7249. <https://doi.org/10.1016/j.mito.2023.01.005>.
17. Kesika P, Thangaleela S, Sisubalan N, Radha A, Sivamaruthi BS, Chaiyasut C. The Role of the Nuclear Factor-Kappa B (NF- κ B) Pathway in SARS-CoV-2 Infection. *Pathogens*. 2024; 13(2):164. <https://doi.org/10.3390/pathogens13020164>.
18. Liu DX, Liang, JQ, Fung, TS Human Coronavirus-229E, -OC43, -NL63, and -HKU1 (Coronaviridae). *Encyclopedia of Virology*, 2021: 428–440. <https://doi.org/10.1016/B978-0-12-809633-8.21501-X>.
19. Mathieu E, Bernard AS, Ching HYV, Somogyi A, Medjoubi K, Fores JR, et al. Anti-inflammatory activity of superoxide dismutase mimics functionalized with cell-penetrating peptides. *Dalton Trans*. 2020 Feb 21;49(7):2323-2330. doi: 10.1039/c9dt04619d.
20. Morgan MJ, Liu ZG. Crosstalk of reactive oxygen species and NF- κ B signaling. *Cell Res*. 2021;21(1):103-115. <https://doi.org/10.3389/fgene.2021.622271>.
21. Muhammad JS, ElGhazali G, Shafarin J, Mohammad MG, Abu-Qiyas A, Hamad M. SARS-CoV-2-induced hypomethylation of the ferritin heavy chain (FTH1) gene underlies serum hyperferritinemia in severe COVID-19 patients. *Biochem Biophys Res Commun*. 2022 Nov 26;631:138-145. doi: 10.1016/j.bbrc.2022.09.083.
22. Muhammad JS, Saheb Sharif-Askari N, Cui ZG, Hamad M, Halwani R. SARS-CoV-2 Infection-Induced Promoter Hypomethylation as an Epigenetic Modulator of Heat Shock Protein A1L (HSPA1L) Gene. *Front Genet*. 2021 Feb 19;12:622271. doi: 10.3389/fgene.2021.622271.
23. Nepogodiev HA, Glasbey D, Li JC, Gujjuri OM, Morton RR, Mantoglu DG. Mortality and pulmonary complications in patients undergoing surgery with perioperative sars-cov-2 infection: An international cohort study. 2020. *The Lancet*, 396 (10243), pp. 27-38. doi: 10.1016/S0140-6736(20)31182-X.
24. Ni W, Yang X, Yang D, Bao J, Li R, Xiao Y, et al. Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19. *Crit Care*. 2020 Jul 13;24(1):422. doi: 10.1186/s13054-020-03120-0.
25. Palmer JE, Wilson N, Son SM, Obrocki P, Wrobel L, Rob M, et al. Autophagy, aging, and age-related neurodegeneration. *Neuron*. 2025 Jan 8;113(1):29-48. doi: 10.1016/j.neuron.2024.09.015.
26. Perreau M, Suffiotti M, Marques-Vidal P. The cytokines HGF and CXCL13 predict the severity and the mortality in COVID-19 patients. *Nat Commun* 12, 4888 (2021). <https://doi.org/10.1038/s41467-021-25191-5>.
27. Saeedi-Boroujeni A, Mahmoudian-Sani MR. Anti-inflammatory potential of Quercetin in COVID-19 treatment. *J Inflamm* 18, 3 (2021). <https://doi.org/10.1186/s12950-021-00268-6>.
28. Şehirli AÖ, Aksoy U, Koca-Ünsal RB, Sayiner S. Role of NLRP3 inflammasome in COVID-19 and periodontitis: Possible protective effect of melatonin. *Med Hypotheses*. 2021 Jun;151:110588. doi: 10.1016/j.mehy.2021.110588.
29. Shi Z, Puyo CA. N-Acetylcysteine to Combat COVID-19: An Evidence Review. *Ther Clin Risk Manag*. 2020;16:1047-1055. Published 2020 Nov 2. doi:10.2147/TCRM.S273700.
30. Talukdar J, Bhadra B, Dattaroy T, Nagle V, Dasgupta S. Potential of natural astaxanthin in alleviating the risk of cytokine storm in COVID-19. *Biomed Pharmacother*. 2020;132:110886. doi:10.1016/j.biopha.2020.110886.
31. VasanthiDharmalingam P, Karuppagounder V, Watanabe K, Karmouty-Quintana H, Palaniyandi SS, Guha A, et al. SARS-CoV-2 Mediated Hyperferritinemia and Cardiac Arrest: Preliminary Insights. *Drug Discov Today*. 2021 May;26(5):1265-1274. doi: 10.1016/j.drudis.2021.01.014.
32. Wang H, Rizvi SRA, Dong D, Lou J, Wang Q, Sopipong W, et al. Emerging variants of SARS-CoV-2 NSP10 highlight strong functional conservation of its binding to two non-structural proteins, NSP14 and NSP16. *Elife*. 2023 Dec 21;12:RP87884. doi: 10.7554/eLife.87884.
33. Weir EK, Thenappan T, Bhargava M, Chen Y. Does vitamin D deficiency increase the severity of COVID-19?. *Clin Med (Lond)*. 2020;20(4):e107-e108. doi:10.7861/clinmed.2020-0301.
34. Wu G. Important roles of dietary taurine, creatine, carnosine, anserine and 4-hydroxyproline in human nutrition and health. *Amino Acids* 52, 329–360 (2020). <https://doi.org/10.1007/s00726-020-02823-6>.

Стаття надійшла 02.02.2024 р.